BURUNG

Burung Kunyit Besar (Dendang Selayang) / Black-Naped Oriole



Nama Saintifik : Oriolus chinensis
Burung jantan spesis ini sepenuhnya berwarna kuning keemasan kecuali bahagian sayap dan ekor serta jalur hitam di bahagian mata. Manakala, Burung betina serupa dengan burung jantan tetapi dengan bahagian atas yang lebih hijau kekuningan. Ketika musim mengawan, persaingan antara sesama burung jantan akan menjadi sengit dengan suasana dipenuhi dengan bunyi burung jantan mempertahankan kawasan mereka serta menarik perhatian burung betina, aktiviti mengejar dan berebut pokok.

The male bird of this species is predominantly golden yellow, except for the wings and tail, and features a black stripe around the eyes. The female bears a resemblance to the male but with a more yellowish-green upperpart. During the mating season, competition among male birds intensifies, with the air filled with their calls as they defend territories and vie for the attention of female birds, engaging in pursuits and skirmishes over trees.

Burung Tiong Batu / Oriental Dollarbird



Nama Saintifik: Eurystomus orientalis
Burung dewasa mempunyai bahagian kepala dan dada berwarna coklat gelap, bahagian kerongkong berwarna biru terang berkilat. Kedua-dua jantina serupa tetapi betina warnanya agak kusam. Ketika musim mengawan, pasangan burung sering dilihat terbang dalam pergerakan berguling disertai dengan bunyi panggilan.

Adult Oriental Dollarbirds feature a dark brown head and chest, contrasted by a vibrant, shiny bright blue throat. Both sexes share similar characteristics, although the female tends to have a duller coloration. Throughout the mating season, pairs of these birds are frequently observed flying in a rolling motion while emitting distinct calls

BURUNG

Burung Pacat Sayap Biru / Blue-winged Pitta



Nama Saintifik: Pitta moluccensis
Burung ini membina sarang yang besar, biasanya di atas tanah, diperbuat daripada ranting, akar, rumput, daun dan lumut. Burung betina bertelur 4-6 telur berwarna putih atau krim dengan tanda ungu, dan kedua-dua ibu bapa bergilir-gilir mengeram telur selama 15-17 hari.

These birds build large nests, usually on the ground, made of twigs, roots, grass, leaves and moss. Females lay 4-6 white or cream eggs with purple markings, and both parents take turns incubating the eggs for 15-17 days.

Burung Pekaka Sungai / Collared Kingfisher



Nama Saintifik: Todiramphus chloris
Burung Pekaka Sungai adalah sejenis burung yang mempunyai bulu pelepah yang berwarna biru terang dan leher berwarna putih. Burung ini boleh berdiam diri bertenggek menunggu mangsa untuk tempoh yang lama. Kedua-dua jantan dan betina mengambil bahagian dalam mengeram telur dan memberi makan kepada anak burung. Burung muda meninggalkan sarang kira-kira 44 hari selepas menetas. Dua ekor anak sering dibesarkan dalam setahun.

The Collared Kingfisher, also known as the White-collared Kingfisher, Black-masked Kingfisher, or Mangrove Kingfisher, is a bird species characterized by bright blue feathers and a white neck. These birds can remain silent while perched, waiting for prey for extended periods. Both parents participate in incubating the eggs and feeding the chicks. The young birds typically leave the nest approximately 44 days after hatching. Often, two broods are raised in a single year.

BURUNG

Burung Takau Rakit / Black and Red Broadbill

Nama Saintifik: Cymbirhynchus macrorhynchos

Takau rakit atau terajan adalah spesis burung takau dalam keluarga Eurylaimidae. Ia spesis tunggal jenis Cymbirhynchus. Pembiakan berlaku seawal dan sepanjang musim kemarau , dengan sarangnya adalah struktur besar dan mudah dilihat yang biasanya tergantung di atas air. Sarang dibina oleh kedua-dua jantina, daripada tumbuhan menjalar, hifa kulat, lumut dan bahan tumbuhan lain.

The black-and-red broadbill is a species of bird in the typical broadbill family, Eurylaimidae. It is the only species in the genus Cymbirhynchus. Breeding occurs early and throughout the dry season, with the nest being a large, conspicuous structure that usually hangs above the water. Nests are built by both sexes, from creeping plants, fungal hyphae, moss and other plant material.

